

# Particles and Universe

## Lecture 3 Properties of particles

Maria Krawczyk, Aleksander F. Żarnecki

Faculty of Physics UW


### Interactions

Masses and lifetimes of elementary particles

Quarks: flavors and color

Conservation laws and quantum numbers:  
baryon number and lepton numbers

# Discoveries – flood of particles in 50-60 of XX



# Zoo of elementary particles

**Definition: elementary particle** - object simpler than nucleus (exception the simplest nucleus H(hydrogen), proton, which is an elementary particle)

- Elementary particles – a lot (~1000) and various (Zoo):  
<http://pdg.lbl.gov/>  
various masses,  
various lifetimes (they can decay !),  
various electric charges,  
various types of interactions,  
various groups (multiplets), etc

Elementary particles can be compound (proton) !  
the simplest particles → **fundamental particles**

# Particles of matter and particles of forces

Today we will focus on matter

# Interactions

## In macro- and micro scales:

- **gravitation** – act between all massive particles, only attraction, responsible for Sun system, large astronomical objects, etc.
- **electromagnetism (e-m, el-mag)** – electric charge of both signs, attraction and repulsion, atoms ...

## In microworld in addition interactions:


- **strong (nuclear)** - bounding nucleons in nuclei (**pions exchange**)  
range  $10^{-15}$  m
- strong fundamental (color)** - between quarks (**gluons exchange**),  
range  $10^{-15}$  m
- **weak** (nuclear), eg. neutron decay, range smaller than for strong  
(pointlike interaction)
- weak fundamental** between quarks and leptons (exchange of  
gauge boson  $W/Z$  ), range  $10^{-18}$  m

# Range of interactions

$$c = \hbar = 1$$

- Interaction in microworld = emission and absorption of bosons (photon, W/Z, gluons..) → exchange of particles
- Range** (Heisenberg, Yukawa) is related to the mass of exchanged particle (carrier of interactions)

$$x \sim 1/M$$



- gravitation and el-mag **infinite range** → graviton mass? photon mass = 0
- color (strong) int. : range ~ **proton radius  $10^{-15}$  m** (although mass of gluons zero, **confinement!**)
- weak int. **range  $10^{-18}$  m**, related to the mass of bosons

$$W/Z \sim 80-90 \text{ GeV}$$

# Strenght of interactions

- gravitation and el-mag are very different - gravitation very weak (gravitation between two protons  $10^{36}$  times weaker than el-mag)
- Strenght's hierarchy at low\* energies:

**strong > electromagn. > weak > gravitation**


*\* low energies: 1 GeV up to 100 GeV*

in the Standard Model – no gravitation!

- Parameter of strenght of elementary action  
→ **coupling constant**

# Fundamental particles in the Standard Model

- Particles like proton  $p$  and neutron  $n$  are bound states of quarks. Fundamental particles of matter : quarks and leptons have no inner structure
- Carriers of interactions:  $W^+W^-$   $Z$ ,  $\gamma$  and gluons - fundamental particles




*March 2015*

- Physics of elementary particles describes the fundamental level up to  $10^{-18}\text{m}$  and energies up to 8 TeV

→ **Standard Model**



# Fundamental particles



# Properties of elementary particles

# Masses


$E = mc^2$ , mass unit = eV/c<sup>2</sup> - we usually neglect c<sup>2</sup> (c=1)

- Neutrino ~ 0 ?  
Electron ~ 0.5 MeV  
Pions (quarks and antiquarks u,d) ~140 MeV  
Proton, neutron (uud, ddu) ~ 1 GeV  
(highest mass ~172 GeV)
- Origin of mass ? – **BEH mechanism! (2012)**
- Is mass of a particle =sum of constituents' masses ?  
Not always - it is **not so** for nucleons, pions  
where it is coming from strong interactions  
 $m = E / c^2$  (masses of quarks negligible)

# Masses of quarks and leptons


## ■ Masses:

|                           |                             |                              |
|---------------------------|-----------------------------|------------------------------|
| <b>u</b>                  | <b>c</b>                    | <b>t</b>                     |
| 3 MeV                     | 1.25 GeV                    | 172 GeV                      |
| <b>d</b>                  | <b>s</b>                    | <b>b</b>                     |
| 7 MeV                     | 150 MeV *                   | 4.5 GeV                      |
| <b><math>\nu_e</math></b> | <b><math>\nu_\mu</math></b> | <b><math>\nu_\tau</math></b> |
| $<5 \cdot 10^{-6}$ MeV    | $<0.27$ MeV                 | $<31$ MeV                    |
| <b>e</b>                  | <b><math>\mu</math></b>     | <b><math>\tau</math></b>     |
| 0.511 MeV                 | 105.7 MeV                   | 1.78 GeV                     |



# Decays

Decay of particles – spontaneous transformation of one particle into others



Eg. **neutron decay** ( $\beta$  decay)

neutron  $\rightarrow$  proton electron plus ?

*(lifetime of neutron  $886\text{ s} = 14,8\text{ min}$ )*


1914 J. Chadwick: in  $\beta$  decay RaB ( $^{214}\text{Pb}$ ) continuous electron energy spectrum - this can not be decay into two particles (energy-momentum conservation)

1927 J. Ellis confirms this finding..

.....

N. Bohr – maybe energy is not conserved?

# Beta decay 1930



This experimental energy spectrum is from G.J. Neary, Proc. Phys. Soc. (London), A175, 71 (1940).

Expected 2-body decay

- **Problem:** nucleus  $(A,Z)$  thought to be  $A$  protons +  $(A-Z)$  electrons
- **Beta decay:**  $(A,Z) \rightarrow (A, Z+1) + e^-$  (two body decay, monoenergetic  $e^-$ )

# Pauli 1930

Offener Brief an die Gruppe der Radioaktiven bei der  
Gesellschafts-Tagung zu Tübingen.

Abschrift

4th December 1930

Dear Radioactive Ladies and Gentlemen,

As the bearer of these lines, to whom I graciously ask you to listen, will explain to you in more detail, how because of the "wrong" statistics of the N and  ${}^6\text{Li}$  nuclei and the continuous beta spectrum, I have hit upon a desperate remedy to save the "exchange theorem" of statistics and the law of conservation of energy. Namely, the possibility that there could exist in the nuclei electrically neutral particles, that I wish to call neutrons, which have spin and obey the exclusion principle and which further differ from light quanta in that they do not travel with the velocity of light. The mass of the neutrons should be of the same order of magnitude as the electron mass (and in any event not larger than 0.01 proton masses). The continuous beta spectrum would then become understandable by the assumption that in beta decay a neutron is emitted in addition to the electron such that the sum of the energies of the neutron and the electron is constant...

From now on, every solution to the issue must be discussed. Thus, dear radioactive people, look and judge. Unfortunately I will not be able to appear in Tübingen personally, because I am indispensable here due to a ball which will take place in Zürich during the night from December 6 to 7....


Your humble servant,

W. Pauli

Pauli also left in his diaries: "Today I have done something which no theoretical physicist should ever do in his life: I have predicted something which shall never be detected experimentally."

## Chadwick discovers neutron (1932):

- Mass of neutron similar to mass of proton: not Pauli's particle!
- **Fermi** introduces name "neutrino" ( $\nu_e$ ), which is different to neutron, and beta decay is decay of neutron:



ANNO IV - VOL. II - N. 12      QUINDICESIMALE      31 DICEMBRE 1933 - XII

### LA RICERCA SCIENTIFICA

ED IL PROGRESSO TECNICO NELL'ECONOMIA NAZIONALE

#### Tentativo di una teoria dell'emissione dei raggi "beta"

Nota del prof. ENRICO FERMI

**Riassunto:** Teoria della emissione dei raggi  $\beta$  delle sostanze radioattive, fondata sull'ipotesi che gli elettroni emessi dai nuclei sono emessi prima della disintegrazione ma vengono formati insieme ad un neutrino, in modo analogo alla formazione di un quante di luce che accompagna un salto quantico di un atomo. Conferma della teoria con l'esperienza.



theory..  
weak int



# Particle decay

Probability to survive (before decaying)

$$P(t) = e^{-t/(\gamma\tau)}$$

$\tau$  = the mean lifetime (**lifetime**) of the particle (at rest)

(reduction *e*-times,  $e = 2,7$ )

(half lifetime if *reduction 2-times*)

$$\gamma = \frac{1}{\sqrt{1 - v^2/c^2}}$$



the Lorentz factor

Decay rate  $\Gamma = 1/\tau$

total, if various decay modes (channels),  $\Gamma = \sum \Gamma_i$

# Lifetime of particles

- Lifetime of the particles – exponential decay

(lifetime  $\rightarrow 1/e$  ( $e=2.7$ ) particles remain)

Universe 13,7 mld years

- Lifetime ( $\tau$ )

- stable particles (the most...):

electron:  $\tau > 4.6 \cdot 10^{26}$  y i proton:  $\tau > 10^{30}$  y

- short lived:  $\sim 10^{-24}$  s




- long lived :  $10^{-6}$  -  $10^{-8}$  s

(eg. mion  $2 \cdot 10^{-6}$  s, charged pions  $2.6 \cdot 10^{-8}$  s)

- **Probablility of decay small - lifetime long and vice versa (strong int.  $\rightarrow$  short lifetime)**

# Production of particles


- In collision of particles two, three, ...  $N$  particles maybe be produced – in agreement with energy and momentum conservation




**CMS at LHC**

# Production of particles

- In collision of particles two, three,... N particles maybe be produced – in agreement with energy and momentum conservation




- However energy of collision can be transfer **totally** into a rest energy of one particle zgodnie z  $E=mc^2$  - **resonance production** :



*discovery of many particles*

# Number of events in e+e- collision



energy of collision (GeV)

Resonances:

collision energy (GeV) = mass of particle

$$\Delta E \Delta t \geq \hbar/2$$

width of resonance line  $\Gamma = 1/\tau$  (in a half of height)

# Types (flavors) quarks

# Ordinary matter and ordinary quarks

Life, <sup>much</sup> of the Universe, <sup>but</sup> not everything

Stable (ordinary) matter

- up-quark (charge  $+2/3$ )
- down-quark (charge  $-1/3$ )
- electron (charge  $-1$ )
- neutrino (no charge and  $\approx$  zero mass)

F. Close

# Ordinary quarks u (up) i d (down)

The most populated quarks  
in the most populated elementary  
particles


**H atom**  
(not to scale!)

**a miracle  
of  
neutrality**

**electron  
balances**

**uud**

hint of unification



from Close



# Mass difference of quarks u and d

- Proton (uud) i neutron (ddu)

Masses:  $m_p=938.3$  MeV ,  $m_n=939.6$  MeV,

$\Delta m=1.3$  MeV

→ difference of mass for quarks d and u;  
other measurements →

mass of u, d – few MeV (*light quarks*)

- Neutron decay (fundamental level) →

**decay of quark d to quark u**

(+ electron + antineutrino el.)

Quark d heavier -> decays to a lighter quark u ..

# Proton is stable, while neutron no, - very good:

- Sun (neutron decay)
- Water exists (proton)

Neutron decay  $n \rightarrow p e^- \bar{\nu}_e$

and

~~$p \rightarrow n e^+ \nu_e$~~

However - why proton is stable...?  
What about other decays?

# Isospin (u,d)

- Heisenberg – 1932 (symmetry of nuclear int.)
- isotopic spin (p,n **dublet**) -2 states of nucleon
- formalism like for spin  $\frac{1}{2}$

- so, isospin  $\frac{1}{2}$  for nucleon  $I = 1/2, I_3 = \pm 1/2$
- for pions - 3 states (**triplet**)

isospin  $I = 1, I_3 = 1, 0, -1$

So,  $p = uud, n = ddu$

$$\pi^+ = u \bar{d} \quad \pi^- = d \bar{u} \quad \pi^0 = \frac{1}{\sqrt{2}} (u \bar{u} - d \bar{d})$$

# Relation

electric charge – isospin?

Electric charge  $Q$

$$Q = I_3 + \frac{1}{2} Y$$

$Y$  - hypercharge

# Strange particle

Strange particles were discovered in cosmic rays  
~ 1950 r

later  
in laboratory  
- first K (kaon)



# Strange particles

- Life time longer than for „similar” particles
- produced in pairs in collision of nucleons

$$\Lambda \rightarrow p + \pi^-$$
$$\tau \approx 3 \cdot 10^{-10} \text{ s}$$

$$\Delta \rightarrow p + \pi^-$$
$$\tau \sim 10^{-23} \text{ s}$$

To describe new „category” of particles  
**S** (strangeness)

# Strangeness

Strange particle  $S \neq 0$ ; observed  $S = 1, 2, 3; -1, -2, -3$

nucleons = proton, neutron  $S=0$

pions  $S=0$

For a system of particles strangeness adds (**additivity**);  
in processes with nucleons, pions it is conserved:

$$S_{\text{initial}} = S_{\text{final}}.$$


On fundamental level: quark s ( $S = -1$ ).

The lightest elementary particle contains one  
strange quark - kaon K (mass 500 MeV):

$K^+ = u \bar{s}$ ,  $K^- = \bar{u} s$ ,  $K^0 = d \bar{s}$ ;  $\Lambda(1116 \text{ MeV}) = uds$

→ mass s quark  $\sim 150 \text{ MeV}$

# Quark Model



In 60-ties XX (multiplets)

Gell-Mann and Zweig:

quark hypothesis

$q \bar{q}$  (mesons)

$qqq$  (baryons)




# Quark diagrams

Octet: mesons  $K$ ,  $\pi$  i  $\eta$


masses  $\sim 500$  MeV

strangeness  $S$



isospin  $I$ :  $-1, -\frac{1}{2}, 0, \frac{1}{2}, 1$


# Octet ( $q \bar{q}$ )




Masses 140- 500 MeV

octet

decouplet




Masy ok 1 GeV



Masy ok 1.7 GeV


# Quarks Model: quarks – fundamental representation of SU(3)\_flavor

quark diagrams



$$3 \times 3 \times 3 = 1 + 8 + 8 + 10$$

$$3 \times \bar{3} = 1 + 8$$



# Heavy flavors

Other phenomena – a need to introduce other flavors

(additive quantum numbers):

1974 (charm)  $C \rightarrow$  quark  $c$

1977 (beauty, bottom)  $B^* \rightarrow$  quark  $b$

1995 (true, top)  $T^* \rightarrow$  quark  $t$

- Particle  $J/\psi$  mass 3 GeV. In collision  $e^+e^-$  very narrow resonans. Why so narrow ? Does contain new type of quarks ? Charmonium  $J/\psi = c \bar{c}$  ( $C=0$ ). There are charmed particles with  $C=\pm 1$ ,  $D^+ = c \bar{d}$ . Mass of  $c$  quark  $\sim 1.5$  GeV (half of mass  $J/\psi$  ).  
<http://capp.iit.edu/hep/bquarkfound.html>
- Particle  $\Upsilon$  (9.5 GeV) - bound state  $b \bar{b}$  ( $B^*=0$ ) -bottonium. Particles with  $B^*\neq 0$  exist . Quark  $b$  mass  $\sim 4.5$  GeV
- Quark mass  $t \sim 172$  GeV  $\rightarrow$  there are no bound state.

## CHARMED MESONS ( $C = \pm 1$ )

$D^+ = c\bar{d}$ ,  $D^0 = c\bar{u}$ ,  $\bar{D}^0 = \bar{c}u$ ,  $D^- = \bar{c}d$ , similarly for  $D^*$ 's

## CHARMED, STRANGE MESONS ( $C = S = \pm 1$ )

## BOTTOM MESONS ( $B = \pm 1$ )

$B^+ = u\bar{b}$ ,  $B^0 = d\bar{b}$ ,  $\bar{B}^0 = \bar{d}b$ ,  $B^- = \bar{u}b$ , similarly for  $B^*$ 's

## BOTTOM, CHARMED MESONS ( $B = C = \pm 1$ )

$B_c^+ = c\bar{b}$ ,  $B_c^- = \bar{c}b$ , similarly for  $B_c^*$ 's

Also BOTTOM, STRANGE MESONS.

# Baryons...

## BOTTOM BARYONS ( $B = -1$ )

$$\Lambda_b^0 = udb, \Xi_b^0 = usb, \Xi_b^- = dsb, \Omega_b^- = ssb$$


etc

# u,d,s + c,b,t

- Multiplets with quarks uds –  
mass of these particles close  
(differences below 1 GeV)
- Multiplets with udsc ?, udscb ?  
differences larger  
symmetry broken...
- Quark t too heavy to make a bound state



# Carriers of flavors are quarks



In this table quarks and leptons are in doublets

Why?


This is dictated by weak interaction eg. in beta decay :  
 $d \rightarrow u,$   
 $\text{neutrino} \rightarrow e$

# Exotics (not $q \bar{q}$ , $qqq$ )

<http://www.wired.com/2014/09/tetraquark-quantum-feud/>

- New type of hadrons
- $X(3872)$
- $Z_c(3900)$
- $Y(4140)$   
four-quark system  
 $c \bar{c} d \bar{u}$   
tetraquark

How to get white hadrons?



# Conservation laws and other quantum numbers

# Energy-momentum conservation

- In any process (collision, decay) a total energy :  
 $\text{energy\_final} = \text{energy\_initial}$

For free particles (3-momentum  $\vec{p}$ ,  $m$ - mass at rest )

$$E^2 = (\vec{p}c)^2 + (mc^2)^2$$

In relativistic theory 4-vector  $(E, \vec{p})$  is conserved ( $c=1$ )

- Particle with mass  $m$  has at rest energy  $E=mc^2$   
and massive particles have more chance to decay
- Energy, momentum and angular momentum are conserved

# Charge conservation

- Electric charge is conserved in Nature  
*that why eg. proton can not decay into electron (plus antyneutrino)*
- Charge of elementary particles – only in definite portions (quanta) → charge quantisation (charge quantum number)
- If in some units el. charge of electron = -1, than el. charge of proton = +1,  
(then for  $u$  quark =  $2/3$ , for  $d$  quark  $-1/3$ !)
- Observed elementary particles have el. charge multiplicative of el. charge of electron –  
so  $n = 0, 1, 2, \dots$  or  $-1, -2$ , ( $n = 0$  neutral particle)

# Baryonic quantum number B

- Proton decay is not forbidden by el. charge conservation, nor energy conservation



Seems that p and leptons are not „relatives”

- New idea: Stückelberg (1938r)  
New quantum number to count nucleons (conserved)
- Tests of this hypothesis:

eg. lack of neutron decay  $\rightarrow e^- e^+$ ?  
(*baryon, z greckiego ciężki*)

New (additive) quantum number: baryonic

proton=+1, neutron=+1 ( $\bar{p}, \bar{n} = -1$ ); baryons  $B \neq 0$

**B is conserved in Nature**

# Hadrons = bound states of quarks

Hadrons

Baryons ( $B \neq 0$ )

3 quarks

Mesons ( $B = 0$ )

quark-antiquarks

*Hadron- gruby, mocny*

np. piony

*mezon - pośredni*

# Quantum numbers of quarks cd.

- Baryonic quantum number  $B$  for  $p$ ,  $n = +1$   
Quarks have  $B = 1/3$ , antiquarks =  $-1/3$
- Electric charge for  $p = +1$  and for  $n = 0$   
quarks  $q = 2/3$  lub  $-1/3$   
antiquarks  $\bar{q} = -2/3$  lub  $1/3$   
np..  $u = 2/3$ ,  $d = -1/3$
- Flavor quantum numbers



# Color – new quantum number

flavor (u,d,s...) – classification/structure of particles

color – dynamics of interaction between quarks

- all quarks in 3 color states
- gluons – double color (color and anticolor (eg. gluon red- antiblue) - 8 states (8 gluons)
- photon „feels” el. charge ( $\rightarrow$  *quantum electrodynamics*), gluon – color (color charge) ( $\rightarrow$  *quantum chromodynamics*)
- Free color charge is not observed , there are no free quarks, they are confined permanently ...

# Confinement of color ?!

Quarks and gluons are colored, but hadrons are „white”. Since they are of types

$$(qqq) \text{ i } (q \bar{q})$$

$N_c=3$  is needed to make these combinations

(hadrons) colorless (*color singlet*)

$$3 \times 3 \times 3 = 1 + 8 + 8 + 10 \quad 3 \times \bar{3} = 1 + 8$$

Confinement – new phenomenon -  
end of ladder:

**molecule** → **atom** → **nucleus** → **nucleon** → **quark?**

**Possibly...**

# Electron quantum number $L_e$

- In many processes electron is accompanied by neutrino (or antineutrino) eg. neutron decay

- **Electron number** (additive): electron  $e = +1$ ,  
electron neutrino  $\nu_e = +1$

- For their antiparticles = -1; other particles = 0
- If electron quantum number is conserved

neutron decay must be:



- „**Crossing reaction**” exists:



Observation of crossing process  $\nu_e p \rightarrow n e^+$

-a discovery of electron (anti)neutrino  $\nu_e$

Cowan, Reines'1956 (Nobel 1995)

Before neutrino – only hypothesis (Pauli 1930)

# Muon and taon quantum number

Massive copies of electronu and el. neutrino  $\nu_e$

muon (1937 - „Who ordered that?” I. Rabi ),

taon (1975, M. Perl, Nobel 1995)

*and their neutrinos*

$\nu_\mu$  1962, Schwartz, L. Lederman, J. Steinberger (Nobel 1988)

$\nu_\tau$  2000

**LEPTONS:** (*lepton* - „lekki”)

electron, mion, taon and their neutrinos

Similarly to  $L_e$  one introduces

muon  $L_\mu$  and taon  $L_\tau$  quantum number

# Lepton quantum number L

Lepton quantum number = sum over individual quantum numbers


$$L = L_e + L_\mu + L_\tau$$

L – conserved in Nature...

# Standard Model

**Quarks (all) :**  
**Baryonic number  $B=1/3$**   
**Leptons (all) :**  
**Leptonic number  $L = 1$**

**Antiquarks  $B = -1/3$**   
**Antileptons  $L = - 1$**



**Individual quantum numbers—**

**electron, muon and taon**