

Particles and Universe

Introduction: early discoveries

Maria Krawczyk, Aleksander Filip Żarnecki



KAPITAŁ LUDZKI
NARODOWA STRATEGIA SPÓJNOŚCI




UNIA EUROPEJSKA
EUROPEJSKI
FUNDUSZ SPOŁECZNY



March 1, 2016

HISTORY OF THE UNIVERSE



t = Time (seconds, years)

E = Energy (GeV)

Key

	quark		neutrino		ion		star
	gluon		bosons		atom		galaxy
	electron		meson		photon		black hole
	muon		baryon				
	tau						

Main goal

Present an up-to-date status of theoretical and experimental **particle and astroparticle physics**, including the most recent achievements of the field.

Follow results coming from experiments at Large Hadron Collider (LHC), laboratory and space based searches for **Dark Matter**, as well as from the neutrino oscillation experiments.

Outline

- Introduction to the Standard Model
- Detection of particles, experiments at large colliders
- Selected aspects of the Theory of Fundamental Interactions
- Tests of the Standard Model
- Search for and study of the properties of the Higgs particle
- Flavour mixing and neutrino oscillations
- Models beyond the Standard Model and search for their signatures
- Origin and evolution of the Universe, Dark matter, Dark energy

- 1 Introduction
- 2 Discovery of electron
- 3 Nature of light
- 4 Radioactivity
- 5 Nucleus and nucleons
- 6 Cosmic rays
- 7 New particles

Electricity and Magnetism

Electromagnetic field influences the motion of a charged object:

$$\frac{d}{dt}m\vec{v} = q\vec{E} + q\vec{v} \times \vec{B}$$

Charges and their motion are the source of the electromagnetic field

Maxwell equations (1865):

$$\begin{aligned}\epsilon_0 \operatorname{div} \vec{E} &= \rho \\ \operatorname{rot} \vec{E} &= -\mu_0 \frac{\partial \vec{B}}{\partial t} \\ \operatorname{div} \vec{B} &= 0 \\ \operatorname{rot} \vec{B} &= \vec{j} + \epsilon_0 \frac{\partial \vec{E}}{\partial t}\end{aligned}$$

Electricity and Magnetism

Electromagnetic field influences the motion of a charged object:

$$\frac{d}{dt}m\vec{v} = q\vec{E} + q\vec{v} \times \vec{B}$$

Charges and their motion are the source of the electromagnetic field

Maxwell equations (1865):

$$\begin{aligned}\epsilon_0 \operatorname{div} \vec{E} &= \rho \\ \operatorname{rot} \vec{E} &= -\mu_0 \frac{\partial \vec{B}}{\partial t} \\ \operatorname{div} \vec{B} &= 0 \\ \operatorname{rot} \vec{B} &= \vec{j} + \epsilon_0 \frac{\partial \vec{E}}{\partial t}\end{aligned}$$

Electric and magnetic fields affect each other

⇒ electromagnetic wave: $c = \frac{1}{\sqrt{\epsilon_0 \mu_0}}$

Electricity and Magnetism

Electromagnetic waves were discovered by Heinrich Hertz in 1886

⇒ great success, validation of the Maxwell theory

But the nature of electricity remained unknown!..

Still, the physics seemed to be an almost a “closed” field,
everything was described...

Electricity and Magnetism

Electromagnetic waves were discovered by Heinrich Hertz in 1886

⇒ great success, validation of the Maxwell theory

But the nature of electricity remained unknown!..

Still, the physics seemed to be an almost a “closed” field,
everything was described...


Except for strange phenomena related to current flow in gases:
spectral lines, cathod and anod rays

⇒ one of main fields of experimental activities


Discovery of X-rays by Wilhelm Röntgen in 1895 was unexpected,
but could still fit in that picture...

Discovery of electron

Joseph Thomson 1897



Thomson studied **cathod rays**





he demonstrated that they bend in the external electric field

He estimated the charge to mass ratio for the electron:

$$\frac{e}{m} \approx 2 \cdot 10^{11} \frac{C}{kg}$$

Discovery of electron

Robert Millikan 1909




By measuring the free fall of tiny olive droplets in air, Millikan estimated the electron charge and, taking the Thomson result on e/m , evaluated **electron mass**:

$$m_e = \frac{1}{1837} \cdot m_H$$

Photoelectric effect

Discovered by Hertz in 1887


In 1902, **Philipp Lenard** demonstrated that the photoelectric effect is observed only for selected **wavelengths** of light:



Photoelectric effect

Discovered by Hertz in 1887

In 1902, **Philipp Lenard** demonstrated that the photoelectric effect is observed only for selected **wavelengths** of light:



This effect could not be explained if light is described as the electromagnetic wave!

Photoelectric effect

In 1905, Albert Einstein proposed a hypothesis that light is a flux of electromagnetic energy quanta, which we call **photons** today.

Photon energy:

$$E_{\gamma} = h\nu = \frac{hc}{\lambda}$$



Photoelectric effect

In 1905, Albert Einstein proposed a hypothesis that light is a flux of electromagnetic energy quanta, which we call **photons** today.

Photon energy:

$$E_{\gamma} = h\nu = \frac{hc}{\lambda}$$



To liberate electron, the photon energy has to be higher than the so called **work function** of the cathod element

⇒ wavelength dependence



Arthur Compton 1923

Photon scattering off electrons



Compton demonstrated that photons carry not only the **energy**, but also the **momentum** \Rightarrow they behave as **particles**

Natural radioactivity

Discovered by Henri Becquerel in 1896.

He tried to verify the hypothesis, that X-ray emission is related to fluorescence - emission of light by a substance (eg. uranium) that has absorbed light or other electromagnetic radiation.



Natural radioactivity


Discovered by Henri Becquerel in 1896.

He tried to verify the hypothesis, that X-ray emission is related to fluorescence - emission of light by a substance (eg. uranium) that has absorbed light or other electromagnetic radiation.



Sun was the brightest light source available at that time.


Unfortunately, the weather was poor in Paris, so Becquerel put the uranium sample and unexposed photographic plate into a drawer...




Radioactivity was intensively studied in the following years.

At the beginning of the XX c. three types of radiation were recognized, based on

penetration of matter



bending in the magnetic field



1903


Nobel prize for H.Becquerel, M.Sklodowska-Curie and P.Curie

1903

Nobel prize for H.Becquerel, M.Sklodowska-Curie and P.Curie

E.Rutherford and F.Soddy published the article explaining radioactivity as a **transformation of elements**: one type of atom emits the radiation while transforming into atom of another element.

- α radiation (Helium nucleus)



- β radiation (electrons)



- γ radiation (energetic photons)




1903

Nobel prize for H.Becquerel, M.Sklodowska-Curie and P.Curie

E.Rutherford and F.Soddy published the article explaining radioactivity as a **transformation of elements**: one type of atom emits the radiation while transforming into atom of another element.

- α radiation (Helium nucleus)



- β radiation (electrons)



- γ radiation (energetic photons)




But the atom was still considered as an elementary object...

1903

Nobel prize for H.Becquerel, M.Sklodowska-Curie and P.Curie

E.Rutherford and F.Soddy published the article explaining radioactivity as a **transformation of elements**: one type of atom emits the radiation while transforming into atom of another element.

- α radiation (Helium nucleus)



- β radiation (electrons)



- γ radiation (energetic photons)





But the atom was still considered as an elementary object...

Strong radioactive sources became one of main tools in research...

Rutherford experiment

Scattering of α particles of gold foil




Scintylation observed when the scattered particle hit the fluorescent screen was used to estimate the scattering angle.

Thomson model:
Uniformly charged atom with point-like electrons moving freely inside ("plum pudding model")


Rutherford experiment

Results of H.Geigera i E.Marsdena measurements (1909):

Expected



Measured




Rutherford model (1911)



Rutherford proposed the model of atom made up of a central charge, surrounded by a cloud of electrons.


The positive charge of the atom (10^{-10} m) is confined to almost point-like (10^{-14} m) **nucleus**



α particles can scatter at much larger angles


After the I World War, Rutherford continued to study nuclei of elements with scattering experiments. He also studied induced nuclear reactions.

In 1919 he reported results proving that the hydrogen nucleus is present in other nuclei \Rightarrow considered as **proton discovery**.



After the I World War, Rutherford continued to study nuclei of elements with scattering experiments. He also studied induced nuclear reactions.

In 1919 he reported results proving that the hydrogen nucleus is present in other nuclei \Rightarrow considered as **proton discovery**.




However, Rutherford still believed that heavy nuclei are build of hydrogen (1H) and helium nuclei (3He and 4He), which he considered elementary.

Discovery of neutron

Existence of neutral constituents in nuclei was already considered by Rutherford in 1920. But there was no experimental evidence.


In 1930, Bothe and Backer discovered that beryllium irradiated with α particles emits radiation which is much more penetrating. This radiation was energetic enough to knock protons out of the paraffin target.



Discovery of neutron

Existence of neutral constituents in nuclei was already considered by Rutherford in 1920. But there was no experimental evidence.


In 1930, Bothe and Backer discovered that beryllium irradiated with α particles emits radiation which is much more penetrating. This radiation was energetic enough to knock protons out of the paraffin target.



In 1932, James Chadwick proved that this was not the gamma radiation, but a new type of particle - neutron.

With discovery of neutron, structure of atomic nuclei became clear.


All radioactive decays observed, could be explained in terms of the element transmutation (change of mass and atomic numbers).




With discovery of neutron, structure of atomic nuclei became clear.

All radioactive decays observed, could be explained in terms of the element transmutation (change of mass and atomic numbers).

The uranium-238 decay chain



Radioactivity



Already Coulomb noticed that charged bodies, even perfectly isolated from the environment, lose their charge with time. **He could not explain that...**

In 1900, Charles Wilson discovered that air molecules are partly ionized. He attributed it to the **natural radioactivity** of Earth elements.


First measurements of Theodor Wulf at the Eiffel Tower showed that ionization does not drop with altitude.

In **1912**, **Victor Hess** showed that ionization is, in fact, increasing at high altitudes.




Detailed measurements conducted by **Victoria Hessa** in **1913** showed that the ionization is increasing fast for altitudes above 2 km.

The only plausible explanation was that it comes from the cosmic space...



Detailed measurements conducted by **Victoria Hessa** in **1913** showed that the ionization is increasing fast for altitudes above 2 km.

The only plausible explanation was that it comes from the cosmic space...




However, it was not clear until 1932, if cosmic rays were charged particles or the energetic gamma radiation...

With radioactive sources energies up to 50 MeV could only be obtained. Cosmic rays were the only source of high energy radiation.

Many experiments were build, using cloud chambers or with photographic emulsion, to search for new phenomena in cosmic ray interactions.

Carl Anderson put his Wilson chamber in a strong **magnetic field** and put a thin **lead plate** in the middle.




With radioactive sources energies up to 50 MeV could only be obtained. Cosmic rays were the only source of high energy radiation.

Many experiments were build, using cloud chambers or with photographic emulsion, to search for new phenomena in cosmic ray interactions.

Carl Anderson put his Wilson chamber in a strong magnetic field and put a thin lead plate in the middle.


In 1932 he observed the particle which had a positive charge (as a proton) but the mass of the same order of magnitude as that of electron.

⇒ positron



The muon was first observed in 1932 by Paul Kunze in Rostock. He observed a particle with positive charge, but the mass seemed to be smaller than that of the proton and higher than that of the electron.

The observation was published, but it was not compatible with the "standard model" of that time...



← muon


← electron

Muon discovery

In 1936 the American physicists Carl D. Anderson and Seth Neddermeyer presented results from the study of a large number of cosmic showers.

They concluded that two components are needed to explain their results:

- non-penetrating, easily stopped by 0.7 to 1.5 cm of lead
 \Rightarrow can be interpreted as “free positive and negative electrons”
- penetrating, losing very little energy in lead
 mass smaller than that of a proton \Rightarrow new particle ?...




Muon discovery

In 1936 the American physicists Carl D. Anderson and Seth Neddermeyer presented results from the study of a large number of cosmic showers.

They concluded that two components are needed to explain their results:

- non-penetrating, easily stopped by 0.7 to 1.5 cm of lead
 ⇒ can be interpreted as “free positive and negative electrons”
- penetrating, losing very little energy in lead
 mass smaller than that of a proton ⇒ new particle ?...




Because of its mass, it was at first thought to be the particle predicted by the Japanese physicist Yukawa Hideki in 1935, to explain the strong force that binds protons and neutrons together in atomic nuclei.

However, it was not interacting strong enough with matter...


In 1947, Cecil Powell's group at Bristol University studied cosmic ray interaction in photographic emulsion.

They discovered that the muons are produced by other particles, pions, which live for only a few hundredths of a microsecond.

One of the reconstructed decay chains:



Pion discovery




Strange particles

In 1948, Powell's team at Bristol University studying cosmic ray interactions with photographic emulsion made another spectacular discovery.

They observed the decay of an unknown particle (coming in from top right) into 3 pions.



The kaon was the first discovered member of the family of "strange" particles...



Primary cosmic rays


Observed in the cosmic space, outside Earth's atmosphere

Composition:

- protons (1H) $\sim 86\%$
- α particles (4He) $\sim 13\%$
- heavier nuclei $\sim 1\%$
- neutrons, electrons $\ll 1\%$

(neglecting neutrinos and gammas)

Same as the “composition of the Universe...”



Cosmic rays

Secondary cosmic rays


Result of the primary cosmic ray interactions in the Earth's atmosphere. Secondary particles, mainly pions and kaons, are copiously produced in these interactions. Both pions and kaons are unstable, produced in their decay chain are muons and electrons.

At the sea level:

- muons $\mu^\pm \sim 70\%$
- electrons $e^\pm \sim 25\%$
- protons and pions $\pi^\pm \sim 3\%$




Average flux:

180 charged particles per $m^2 \cdot s$



Production and decay of a Λ particle:

Lifetime much longer
than expected



Strange particles

Production and decay of a Λ particle:


Lifetime much longer
 than expected

$$\Lambda \rightarrow p + \pi^-$$

$$\tau \approx 3 \cdot 10^{-10} \text{ s}$$

$$\Delta \rightarrow p + \pi^-$$

$$\tau \sim 10^{-23} \text{ s}$$






Strange particles

Many new particles discovered after 1950.

They could be grouped into multiplets, based on their properties.

It started to remind the periodic table of elements.

⇒ indication of a more fundamental description.



Concurrently to the scientific discoveries new research tools were being developed

1897 – electron

1911 – atomic nucleus cloud chamber

1919 – proton

1929 – Van der Graaff accelerator

1931 – neutron Lawrence cyclotron

1932 – positron Cockcroft-Walton generator

1937 – muon μ^\pm

Concurrently to the scientific discoveries new research tools were being developed

1897 – electron

1911 – atomic nucleus cloud chamber

1919 – proton

1929 – Van der Graaff accelerator

1931 – neutron Lawrence cyclotron

1932 – positron Cockcroft-Walton generator

1937 – muon μ^\pm

1947 – pion π^\pm

1949 – kaon K^\pm

1952 – bubble chamber

1955 – synchrotron

⇒ dawn of modern particle physics

Changing landscape

