

Particles and Universe: Tests of the SM and Higgs bozon discovery

Maria Krawczyk, Aleksander Filip Żarnecki



KAPITAŁ LUDZKI
NARODOWA STRATEGIA SPÓJNOŚCI



UNIA EUROPEJSKA
EUROPEJSKI
FUNDUSZ SPOŁECZNY



May 17, 2016

- 1 Introduction
- 2 W^\pm and Z^0 bosons
- 3 Top quark
- 4 Higgs boson
- 5 Tests of the SM

Nobel Prizes

- 1979** - Sheldon L. **Glashow**, Abdus **Salam** i Steven **Weinberg**
model of electro-weak interactions, predicting W^\pm and Z^0
- 1984** - Carlo **Rubia** and Simon **Van der Meer**
 W^\pm and Z^0 discovery at CERN SPS
- 1999** - Gerardus 't **Hooft** and Martinus J.G. **Veltman**
renormalization of the Standard Model
- 2004** - David J. **Gross**, H. David **Politzer** and Frank **Wilczek**
model of strong interactions
- 2008** - Yoichiro **Nambu**, Makoto **Kobayashi** and Toshihide **Maskawa**
mechanism of the **spontaneous symmetry breaking**
description of **quark mixing**, predicting 3rd quark generation
- 2013** - Francois **Englert**, Peter **Higgs**
for the mechanism explaining the origin of particle masses
confirmed recently by the ATLAS and CMS experiments


Standard Model: electro-weak int. + **quantum chromodynamics** (QCD)

W^\pm and Z^0 bosons

Weinberg-Salam model (1968)

New model of the weak interactions
Interaction can be described by the exchange of the **very massive** boson: W^\pm or Z^0 .

Muon decay:



“Weakness” is not due to the small coupling but results from the **large boson mass**:

$$G_F \sim \frac{g^2}{m_W^2}$$

Assuming **coupling** g same as for the EM interactions **Weinberg i Salam predicted** masses of W^\pm and Z^0 :

$$m_W \sim 80 \text{ GeV}$$


$$m_Z \sim 90 \text{ GeV}$$

Neutrino interactions with Z^0 (**neutral currents**) exchange - 1973

Discovery SPS accelerator at CERN

In $p\bar{p}$ interactions, $q\bar{q}$ annihilating in virtual photon can result in lepton-pair production (e^+e^- , $\mu^+\mu^-$, $\tau^+\tau^-$).


Drela-Yana process:




If the invariant mass of annihilating quarks is large, also virtual Z^0 can be exchanged.

Z^0 contribution \Rightarrow maximum in the lepton **invariant mass** distribution

UA1 results (1983):



Z^0 event in UA1 detector (1983)



W^\pm and Z^0 bosons

Discovery

In $p\bar{p}$ interactions, $q\bar{q}'$ annihilation can also result in production of the W^\pm boson:

$$u\bar{d} \rightarrow W^+ \rightarrow e^+ \nu_e$$

Decay with **neutrino production**

⇒ transverse momentum imbalance

Longitudinal neutrino momentum unknown

⇒ we can not reconstruct W^\pm mass directly

“Transverse mass” m_T :

reconstructed assuming $p_z^\nu = 0 \Rightarrow m_T \leq m_W$

W^\pm and Z^0 bosons were discovered at UA1 and UA2 experiments at CERN SPS.

UA1 results (1983):




Fig. 3b

W^- event in UA1 detector (1983)



LEP, CERN, Geneva

1989 - 2000: precise tests of the Standard Model



W^\pm and Z^0 bosons




High precision tests of the Standard Model were possible in e^+e^- interactions at LEP and SLC (millions of events).

Clear maximum in the hadron production cross section corresponds to the real Z^0 production (on mass shell).


Width of the maximum corresponds to the finite Z^0 width (Heisenberg's uncertainty principle)

As the Z^0 lifetime is extremely short, its mass is not fixed - and can vary from event to event. Resonance width $\Gamma_Z \approx 2.5$ GeV




$$e^+e^- \rightarrow W^+W^-$$

At LEP W^\pm can only be produced in pairs.
 Three diagrams contribute:



Couplings are uniquely given by the structure of the **Standard Model**
 \Rightarrow strict theoretical prediction



Cancellation of divergences confirmed by LEP experiments

Short history

- 1964 - Gell-Mann and Zweig, quark model with 3 quarks: u, d, s
- 1970 - Glashow, Iliopoulos and Maiani (GIM) - 2 doublets: u, d, s, c
- 1973 - Kobayashi and Maskawa add 3rd doublet (t and b),
to allow for CP violation
- 1974 - Ting, Richter: charm discovery (c quark)
- 1977 - Lederman (Fermilab): b quark discovery

b quark properties (charge, isospin, gauge boson couplings) in agreement with expectation for “down” member of quark doublet

⇒ “up” partner needed: **top**

First prediction (rule of “3”):

$$m_s = 0.5 \text{ GeV} \quad m_c = 1.5 \text{ GeV} \quad m_b = 4.5 \text{ GeV} \quad \Rightarrow \quad m_t \sim 15 \text{ GeV}$$

First “discovery”: CERN SPS, 1984, $m_t \sim 40 \text{ GeV}$ (revoked)

Searches at LEP and HERA, without success...

Precise measurements at LEP

Comparison


Many different observables measured with high precision at LEP.

Only **three free parameters** describing Standard Model interactions (+ fermion and Higgs masses).

Possible choice: α_{em} , G_F , M_Z can be fitted to precise data.

All other electroweak measurements should be then uniquely predicted

$$\text{Level of agreement: } \text{pull} \equiv \frac{|X_{meas} - X_{SM}|}{\sigma_X} \Rightarrow$$



March 2012


Precise measurements at LEP

Corrections

Measurements at LEP turn out to be very sensitive to “higher order” corrections.

In particular, sizable corrections are due to diagrams including exchange of heavy virtual particles: W^\pm boson, top quark, Higgs boson or new exotic states...


Precise measurements at LEP and other experiments allow us to infer about masses of these heavy states, even if we do not observe them directly!



Corrections

We can extract masses of heavy particles from the precision measurements at lower energies.

It worked very well for $W^\pm \Rightarrow$




March 2012

Direct measurement in agreement (to 0.03%) with theoretical predictions based on lower energy measurements.


Top quark

Prediction

Based on the LEP1/SLD measurements, we expected that mass of the top quark should be about 120-180 GeV.




Final LEP1+SLD constraints:




Top quark

Tevatron $p\bar{p}$ interactions at $\sqrt{s} \sim 2 \text{ TeV}$

Top quark pairs produced mainly in quark-antiquark annihilation:



Top decays almost immediately, (no bound state is formed):




We expect to observe a b quark jet plus lepton and neutrino (missing transverse momentum) or pair of jets with invariant mass of W

Top quark

Tevatron


As top is produced in pairs ($t\bar{t}$), 3 event topologies are possible:

leptonic




“gold sample” but only
~4% of events
(e^+e^- , $\mu^+\mu^-$ i $e^\pm\mu^\pm$)

semi-leptonic



~30% of events
(e^\pm and μ^\pm only)
small background

hadronic



~46% of events
huge background
difficult identification

Top quark

Tevatron

One of the first events




$e + 4$ jet event

40758_44414
24-September, 1992

TWO jets tagged by SVX


fit top mass is $175 \pm 10 \text{ GeV}/c^2$

e^+ , Missing E_T , jet #4 from top
jets 1,2,3 from top (2&3 from W)



⇒ actual discovery in 1995

Number of jets reconstructed in events with isolated lepton (e^\pm, μ^\pm)



Events with lepton and ≥ 3 jets

⇒ mostly $t\bar{t}$



Thousands of events collected...

Top quark

Tevatron

Distribution of the reconstructed top mass in one of the considered channels



Comparison of results



CDF + D0:
 $m_t = 174.34 \pm 0.64$ GeV

Comparison


Direct measurements compared with constraints obtained with precise measurements at lower energies.



Direct measurement in agreement with theory predictions based on precise measurements at lower energies.

LHC experiments



High precision studies of processes with W^\pm , Z^0 and top quark production.
 All results with perfect agreement with SM predictions. **March 2014:**



Top quark

LHC experiments

High precision studies of processes with W^\pm , Z^0 and top quark production. All results with perfect agreement with SM predictions.




Number of t produced at LHC is already much higher than at Tevatron...

Corrections

We can extract **masses of heavy particles** from the **precision measurements** at **lower energies**.

For W^\pm and t direct measurements in perfect agreement with expectation

⇒ we can make another step and try to estimate mass of the yet unobserved particle - **Higgs**



Higgs mass

Analysis of **all** available **data** indicated, that the Higgs mass should be **about 100 – 200 GeV**

2010 fit result:

$$m_h = 89^{+35}_{-26} \text{ GeV}$$


or: $m_h < 158 \text{ GeV}$ (95% CL)

Limit from direct search:

$m_h > 114.4 \text{ GeV}$ (95% CL)

all LEP data:

ALEPH + DELPHI + L3 + OPAL




LHC, CERN, Genewa

Run I: 2009-2012 Run II: 2015-2018(?)



Higgs at LHC

Higgs boson is a key element of the Standard Model, with very **special properties**. Search for the Higgs boson and measurement of its properties were the main goals of LHC experiments.



Higgs boson decays primarily to the **heaviest available** states: for mass $m_h < 135\text{ GeV}$ decay to $b\bar{b}$ dominates.

However, there is a huge background of other pp collisions with $b\bar{b}$ production.


We need to look for decay channels with less background...

Higgs at LHC

For small Higgs masses the best channel is

$$H \rightarrow \gamma\gamma$$

Background is high, but we should see a **clear Higgs peak**




For large masses we should look for:

$$H \rightarrow Z^0 Z^0 \rightarrow l^+ l^- l^+ l^-$$

as the charged leptons (e^\pm ; μ^\pm) are very **easy to identify**.


But the events are rare...



computer simulation results

Higgs at LHC


$H \rightarrow \gamma\gamma$



ATLAS EXPERIMENT

Run Number: 191426, Event Number: 86694500

Date: 2011-10-22 15:30:29 UTC






CMS Experiment at the LHC, CERN

Data recorded: 2011-01-25 06:34:20 683895 647061410.01977

Run / Event: 167075 / 6168850867



$H \rightarrow Z^0 Z^0 \rightarrow e^+ e^- e^+ e^-$



Higgs at LHC

In December 2011, **ATLAS** and **CMS** experiments presented first results of the Higgs boson search, based on the data sample collected in 2010-2011.



Event statistics still very low



Higgs at LHC

In December 2011, **ATLAS** and **CMS** experiments presented first results of the Higgs boson search, based on the data sample collected in 2010-2011. **Event statistics still very low**

$$H \rightarrow Z^0 Z^0 \rightarrow l^+ l^- l^+ l^-$$



Higgs mass

Analysis of **all** available **data** from LEP, Tevatron and the first data from LHC (!).


In spring 2012 only a small gap remained where the Higgs mass could be in agreement with the Standard Model predictions:

$$114.7 \text{ GeV} < m_H < 127 \text{ GeV}$$

If this “window” is closed, it would mean that the Standard Model is **excluded!**



But the measurement most difficult for mass of about 120 GeV...

Intermediate mass region, between “light” and “heavy” Higgs



Results of **ATLAS** and **CMS**, after including data collected in 2012.



Fivefold increase in event sample! **Signal clearly visible**



Higgs at LHC

Results of **ATLAS** and **CMS**, after including data collected in 2012.
Fivefold increase in event sample! **Signal clearly visible**

$$H \rightarrow Z^0 Z^0 \rightarrow l^+ l^- l^+ l^-$$




Higgs at LHC

Results of **ATLAS** and **CMS**, after including data collected in 2012.

Two considered “discovery channels” also give the **most precise** mass determination for new particle.



Results based on the full 2009-2012 LHC data sample (Run I):

$$M_H = 125.09 \pm 0.24 \text{ GeV}$$



Large sample of collected events allowed searching for other decay channels

Number of events for all identified production and decay channels in good agreement with SM



Only 2/3 of measurements expected to agree with predictions within $\pm 1\sigma$

Higgs couplings


Key prediction of the model can now be verified:

Higgs boson coupling should be proportional to the particle mass

Errors are still very large!

Much higher precision possible after more data collected at $\sqrt{s}=13\text{TeV}$.

Tenfold increase in data sample expected until 2020.




Higgs at LHC

Standard Model

Within the Standard Model, Higgs boson mass is related to W and t masses via loop corrections.



Higgs boson mass measurement at LHC in agreement with direct measurements of W and t masses, and LEP results.

Everything looks fine...



First look at Run 2 results

$$H \rightarrow ZZ^* \rightarrow 4l$$




Not much luminosity delivered in 2015. Waiting for more data...

SM tests

LHC results


High precision studies of processes with W^\pm , Z^0 and H boson, and top quark production. Perfect agreement with SM predictions. **Nov. 2015:**



SM tests

LHC results



High precision studies of processes with W^\pm , Z^0 and H boson, and top quark production. Perfect agreement with SM predictions. **Nov. 2015:**



SM tests

LHC results

In 2015 LHC was running at $\sqrt{s} = 13$ TeV, little data collected.
Small excess observed in two photon production $pp \rightarrow \gamma\gamma + X$:



$$m_{\gamma\gamma} = 750 \text{ GeV} \Rightarrow 3.6 \sigma \text{ (} 2.0\sigma \text{)}$$

$$m_{\gamma\gamma} = 760 \text{ GeV} \Rightarrow 2.6 \sigma \text{ (} 1.2\sigma \text{)}$$

Not yet convincing...

LHC project spans over many decades

- 1984: first LHC project draft
- 1992: proposals for ATLAS and CMS experiments
- 1989: LEP started
- 1994: LHC construction approved, 1997: ATLAS and CMS approved
- 2008: LHC fails to start - magnet explosion
- 2009: successful start of LHC
- 2010-2012: pp collisions at $\sqrt{s}=7-8$ TeV (Run 1, $\sim 25fb^{-1}$)
 - 2013-2014: Long Shutdown 1 (LS1) - magnet protection upgrade
- 2015-2018: pp collisions at $\sqrt{s}=13-14$ TeV (Run 2, $\sim 100fb^{-1}$)
 - 2019-2020: Long Shutdown 2 (LS2) - LHC/experiment upgrades
- 2021-2023: pp collisions at $\sqrt{s}=14$ TeV (Run 3, $\sim 200fb^{-1}$)
 - 2024-2026: Long Shutdown 3 (LS2) - LHC/experiment upgrades
- 2026-2035: pp collisions at $\sqrt{s}=14$ TeV (HL-LHC, $\sim 3000fb^{-1}$)

Summary

A new era in particle physics started in 2012.

ATLAS and **CMS** experiment discovered new particle, with properties consistent with that of the Higgs boson (50 year after its prediction)

Great success of the Standard Model!

and thousands of researchers, engineers, technicians and students involved for over 20 years in preparations of the LHC experiments.

All LHC results in good agreement with SM so far!

But we also face new challenges now:

- we need to precisely measure properties of the new particle
- we have to keep looking for other new objects
eg. “dark matter” particles

We do believe there is “something” beyond the Standard Model.
Higgs is just the begining...