

CLIC sensitivity to dark matter production with light mediator exchange

Jan Kalinowski^a, Wojciech Kotlarski^b, Krzysztof Mekala^a, Pawel Sopicki^a,
Aleksander Filip Żarnecki^a



^a Faculty of Physics, University of Warsaw

^b Institut für Kern- und Teilchenphysik, TU Dresden


Research supported by  NATIONAL SCIENCE CENTRE
POLAND

CLICdp WG Analysis Meeting
February 8, 2021

- 1 Motivation
- 2 Simulating mono-photon events
- 3 Results
- 4 Plans & Conclusions

Dark Matter production

The mono-photon signature is considered to be the most general way to look for DM particle production in future e^+e^- colliders.



DM can be pair produced in the e^+e^- collisions via exchange of a new mediator particle, which couples to both electrons (SM) and DM states


This process can be detected if additional hard photon radiation from the initial state is observed in the detector...

New analysis approach

Most of the studies performed so far focused on heavy mediator exchange (EFT limit) and coupling values $\mathcal{O}(1)$
 \Rightarrow extracted were limits on DM or mediator masses

In our study:

- focus on light mediator exchange (DM even lighter)
- consider very small mediator coupling to SM, $\Gamma_{SM} \ll \Gamma_{tot}$



From [arXiv:1910.11775](https://arxiv.org/abs/1910.11775)
 ESPP Physics Briefing Book


“Experimental-like” approach \Rightarrow focus on cross section limits

New analysis approach

Most of the studies performed so far focused on heavy mediator exchange (EFT limit) and coupling values $\mathcal{O}(1)$
 \Rightarrow extracted were limits on DM or mediator masses

In our study:

- focus on light mediator exchange (DM even lighter)
- consider very small mediator coupling to SM, $\Gamma_{SM} \ll \Gamma_{tot}$



ILD study: [arXiv:2001.03011](https://arxiv.org/abs/2001.03011)
 Phys. Rev. D 101, 075053 (2020)

“Experimental-like” approach \Rightarrow focus on cross section limits


Dedicated simulation procedure with Whizard

Two variables, calculated separately for each emitted photon:

$$q_- = \sqrt{4E_0 E_\gamma} \cdot \sin \frac{\theta_\gamma}{2},$$

$$q_+ = \sqrt{4E_0 E_\gamma} \cdot \cos \frac{\theta_\gamma}{2},$$



are used to separate “soft ISR” emission region from the region described by ME calculations.



All “detectable” photons are simulated with Matrix Elements

Validation of the procedure

WHIZARD predictions were compared to the results from the KKMC code for $e^+e^- \rightarrow \nu\bar{\nu} + N\gamma$





⇒ very good agreement observed (both for shape and normalisation)

For more details:

J. Kalinowski et al., Eur. Phys. J. C 80 (2020) 634, arXiv:2004.14486

Detector simulation

CLICdet model for Delphes modified to include forward calorimeters



Included in the official Delphes repository as
`delphes_card_CLICdet_Stage3_fcal.tcl`

Event selection

On generator level:

- 1, 2 or 3 ME photons
nonradiative events for signal only (for normalisation)
- all ME photons with $q_{\pm} > 1 \text{ GeV}$ & $E^{\gamma} > 1 \text{ GeV}$
rejected are events with $q_{\pm} > 1 \text{ GeV}$ & $E^{\gamma} > 1 \text{ GeV}$ for any of the ISR photons
- at least one ME photon with $p_T^{\gamma} > 5 \text{ GeV}$ & $7^{\circ} < \theta^{\gamma} < 173^{\circ}$

On detector simulation level:


- single photon with $p_T^{\gamma} > 10 \text{ GeV}$ & $|\eta^{\gamma}| < 2.6$
- no other activity in the detector
other reconstructed objects
 - no electrons
 - no LumiCal photons
 - no BeamCal photons
 - no jets ($p_T > 20 \text{ GeV}$)

Background distributions


for CLIC running at 3 TeV

Two SM backgrounds considered:

Bhabha scattering and (radiative) neutrino pair production



negative e^- polarisation
4000 fb^{-1}




positive e^- polarisation (80%)
1000 fb^{-1}

Background distributions


for CLIC running at 3 TeV

For mono-photon events, two variables fully describe event kinematics

⇒ use 2D distribution of $(\log_{10} E_\gamma, \eta)$ to constrain DM production



negative e^- polarisation
 4000 fb^{-1}



positive e^- polarisation (80%)
 1000 fb^{-1}

Simplified DM model

Simplified model covering most popular scenarios of DM pair-production

Possible DM candidates:


- real or complex scalar
- Majorana or Dirac fermion
- real vector

Possible mediators:

- scalar
- pseudo-scalar
- vector
- axial-vector

(mixed couplings, eg. V-A or V+A, also possible)

Cross section for $e^+e^- \rightarrow \chi\chi$ for
 $M_\chi = 50 \text{ GeV}$ and $M_Y = 300 \text{ GeV}$



Simplified DM model

Simplified model covering most popular scenarios of DM pair-production

Possible DM candidates:


- real or complex scalar
- Majorana or Dirac fermion
- real vector

Possible mediators:

- scalar
- pseudo-scalar
- vector
- axial-vector

(mixed couplings, eg. V-A or V+A, also possible)

Cross section for $e^+e^- \rightarrow \chi\chi$ at
3 TeV CLIC for $M_\chi = 50$ GeV



Simplified DM model

Simplified model covering most popular scenarios of DM pair-production

Possible DM candidates:


- real or complex scalar
- Majorana or Dirac fermion
- real vector

Possible mediators:

- scalar
- pseudo-scalar
- vector
- axial-vector

(mixed couplings, eg. V-A or V+A, also possible)

Cross section for $e^+e^- \rightarrow \chi\chi$ at
500 GeV ILC for $M_\chi = 50$ GeV




Tagging efficiency


Detectable hard photon emitted only in a fraction of signal event

$$\sigma(e^+e^- \rightarrow \chi\chi\gamma_{\text{tag}}) = f_{\text{mono-photon}} \cdot \sigma(e^+e^- \rightarrow \chi\chi(\gamma))$$

ILC @ 500 GeV



CLIC @ 3 TeV




Emission strongly suppressed for narrow mediator with $M_Y \sim \sqrt{s}$

Signal distributions for CLIC at 3 TeV


For fermion DM with $M_\chi = 50$ GeV and vector mediator with $\Gamma/M = 0.03$

Mediator mass: $M_\gamma = 400$ GeV

Signal for negative polarisation



Signal for positive polarisation




Signal normalised to the DM pair-production cross section of 1 fb

Signal distributions for CLIC at 3 TeV


For fermion DM with $M_\chi = 50$ GeV and vector mediator with $\Gamma/M = 0.03$

Mediator mass: $M_\gamma = 1200$ GeV

Signal for negative polarisation



Signal for positive polarisation




Signal normalised to the DM pair-production cross section of 1 fb

Signal distributions for CLIC at 3 TeV


For fermion DM with $M_\chi = 50$ GeV and vector mediator with $\Gamma/M = 0.03$

Mediator mass: $M_\gamma = 2400$ GeV

Signal for negative polarisation



Signal for positive polarisation




Signal normalised to the DM pair-production cross section of 1 fb

Signal distributions for CLIC at 3 TeV


For fermion DM with $M_\chi = 50$ GeV and vector mediator with $\Gamma/M = 0.03$

Mediator mass: $M_\gamma = 2800$ GeV

Signal for negative polarisation



Signal for positive polarisation




Signal normalised to the DM pair-production cross section of 1 fb

Signal distributions for CLIC at 3 TeV


For fermion DM with $M_\chi = 50$ GeV and vector mediator with $\Gamma/M = 0.03$

Mediator mass: $M_\gamma = 3200$ GeV

Signal for negative polarisation



Signal for positive polarisation




Signal normalised to the DM pair-production cross section of 1 fb

Signal distributions for CLIC at 3 TeV


For fermion DM with $M_\chi = 50$ GeV and vector mediator with $\Gamma/M = 0.03$

Mediator mass: $M_\gamma = 12000$ GeV

Signal for negative polarisation




Signal for positive polarisation




Signal normalised to the DM pair-production cross section of 1 fb

Cross section limits for radiative events at CLIC (tagged photon)

Vector mediator with $\Gamma/m = 3\%$




Combined limits for vector mediator




Limits calculated with CL_s approach using RooFit v3.60

Cross section limits for radiative events at CLIC (tagged photon)

Vector mediator with $\Gamma/m = 3\%$




Combined limits for vector mediator




Limits calculated with CL_s approach using RooFit v3.60

Cross section limits for radiative events at CLIC (tagged photon)

Vector mediator with $\Gamma/m = 3\%$




Combined limits for $\Gamma/m = 3\%$




Limits calculated with CL_s approach using RooFit v3.60

Cross section limits for total DM production cross section at CLIC

Vector mediator with $\Gamma/m = 3\%$




Combined limits for vector mediator




4000 fb^{-1} for -80% e^- polarisation + 1000 fb^{-1} for $+80\%$ e^- polarisation

Cross section limits for total DM production cross section at ILC

Vector mediator with $\Gamma/m = 3\%$



Combined limits for vector mediator





1600 fb^{-1} for both $-80\%/+30\%$ and $+80\%/-30\%$ e^-/e^+ polarisation

Coupling limits at CLIC

Combined coupling limits for assumed mass and width of the mediator.

For vector mediator




Almost uniform sensitivity to g_{eeY} up to kinematic limit.


Coupling limits at CLIC

Combined coupling limits for assumed mass and width of the mediator.

For vector mediator



Model comparison





Coupling limits almost independent on the assumed coupling structure!

Coupling limits at ILC

Combined coupling limits for assumed mass and width of the mediator.


For vector mediator




SM Branching Ratio limits

Branching Ratio for mediator decay to e^+e^-
 corresponding to the extracted cross section and coupling limits

CLIC @ 3TeV



ILC @ 500 GeV



For $M_Y < \sqrt{s}$ numbers of expected events too small
 to search for resonant mediator production in e^+e^- decay channel!


Effective mass scale limits

$$\Lambda^2 = \frac{M_Y^2}{|g_{eeY}g_{\chi\chi Y}|}$$


For $M_Y \gg \sqrt{s}$, limits on the effective mass scale of new interactions no longer depend on the assumed mediator mass or width

⇒ EFT approximation can be used

CLIC @ 3TeV



ILC @ 500 GeV



Summary

Analysis of mono-photon events at 3 TeV sensitive to wide range of DM pair-production scenarios

- $\mathcal{O}(1 \text{ fb})$ limits on the radiative production cross section

$$e^+e^- \rightarrow \chi\chi\gamma^{\text{tag}}$$

- $\mathcal{O}(10 \text{ fb})$ limits on the DM pair-production

$$e^+e^- \rightarrow \chi\chi(\gamma)$$

except for the resonance region $M_\gamma \sim \sqrt{s}$

- $\mathcal{O}(10^{-2})$ limits on the mediator coupling to electrons

up to the kinematic limit $M_\gamma \leq \sqrt{s}$

- mono-photon analysis limits more stringent than the limits expected from direct resonance search in SM decay channels
- for heavy mediators, limits from EFT analysis can be reproduced

Plans

For the next weeks:

- verify stability of the results: influence of cuts, binning etc.
- include systematic uncertainties
- prepare results for LaThuile'2021 and LCWS'2021

Future options:

- compare more DM production scenarios
- estimate discovery range
- consider mediator mass and width determination
- estimate limits expected from Bhabha measurement

Conclusions

New framework for **mono-photon analysis** developed

- different scenarios possible with simplified DM model
- focus on light mediator exchange
- consider very small mediator coupling to SM, $\Gamma_{SM} \ll \Gamma_{tot}$

Mono-photon analysis at 3 TeV CLIC sensitive to **light mediators** for SM **coupling values down to $\mathcal{O}(10^{-2})$** :

- comparable sensitivity of negative and positive polarisation samples **4000 fb^{-1} and 1000 fb^{-1} , respectively**
- limits largely independent on the mediator type/coupling
- limits stronger than the estimated sensitivity from the direct resonance search

Thank you!

Simplified DM model

Dark matter particles, X_i , couple to the SM particles via an mediator, Y_j .

Each simplified scenario is characterized by **one dark matter candidate** and **one mediator** from the set listed below:

	particle	mass	spin	charge	self-conjugate	type
DM	X_R	m_{X_R}	0	0	yes	real scalar
	X_C	m_{X_C}	0	0	no	complex scalar
	X_M	m_{X_M}	$\frac{1}{2}$	0	yes	Majorana fermion
	X_D	m_{X_D}	$\frac{1}{2}$	0	no	Dirac fermion
	X_V	m_{X_V}	1	0	yes	real vector
mediator	Y_R	m_{Y_R}	0	0	yes	real scalar
	Y_V	m_{Y_C}	1	0	yes	real vector
	T_C	m_{T_C}	0	1	no	charged scalar

Simplified DM model

Lagrangian describing mediator coupling to electrons given by

$$\mathcal{L}_{eeY} \ni \bar{e}(g_{eY_R}^1 + v\gamma^5 g_{eY_R}^5)eY_R + \bar{e}\gamma_\mu(g_{eY_V}^1 + \gamma^5 g_{eY_V}^5)eY_V^\mu$$

The interaction of mediators with dark matter is described by

$$\begin{aligned} \mathcal{L}_{XXY} \ni & g_{X_R Y_R} X_R^2 Y_R + ig_{X_C Y_V}(X_C^*(\partial_\mu X_C) - (\partial_\mu X_C^*)X_C)Y_V^\mu + \\ & \bar{X}_D(g_{X_D Y_R}^1 + v\gamma^5 g_{X_D Y_R}^5)X_D Y_R + \bar{X}_D\gamma_\mu(g_{X_D Y_V}^1 + \gamma^5 g_{X_D Y_V}^5)X_D Y_V^\mu \\ & \bar{X}_M(g_{X_M Y_R}^1 + v\gamma^5 g_{X_M Y_R}^5)X_M Y_R + g_{X_M Y_V}^5 \bar{\psi}_M \gamma_\mu \gamma^5 \psi_M Y_V^\mu \end{aligned}$$


ILC vs CLIC comparison of simulation and analysis setup

	ILCgen	CLICdet
Detector acceptance (Delphes model)		
tracking	$ \eta < 3$	$ \eta < 2.54$
ECAL	$ \eta < 3$	$ \eta < 3$
LumiCal	$3 < \eta < 4$	$3 < \eta < 4$
BeamCal	$4 < \eta < 5.8$	$4 < \eta < 5.3$
Generator level cuts		
p_T^γ min.	2 GeV	5 GeV
Θ^γ min.	5°	7°
Detector level cuts		
p_T^γ min.	3 GeV	10 GeV
$ \eta^\gamma $ max.	2.8 (7°)	2.6 (8.5°)


ISR rejection efficiency

Fraction of events generated by WHIZARD removed by ISR rejection procedure (ISR photons emitted in the phase-space region covered by ME)

ILC @ 500 GeV



CLIC @ 3 TeV



Cross section limits for radiative events at 3 TeV CLIC

Vector mediator with $\Gamma/m = 1\%$

